



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Molecular signatures for low temperature memory in Arabidopsis

Zuther, Ellen; Schaarschmidt, Stephanie; Fischer, Axel; Erban, Alexander; Pagter, Majken; Mubeen, Umarah ; Walther, Dirk ; Giavalisco, Patrick ; Kopka, Joachim ; Sprenger, Heike; Hinch, Dirk K.

Publication date:
2019

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Zuther, E., Schaarschmidt, S., Fischer, A., Erban, A., Pagter, M., Mubeen, U., Walther, D., Giavalisco, P., Kopka, J., Sprenger, H., & Hinch, D. K. (2019). *Molecular signatures for low temperature memory in Arabidopsis*. Abstract from SEB's Annual Meeting, Sevilla, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Molecular signatures for low temperature memory in *Arabidopsis*

Ellen Zuther¹, Stephanie Schaarschmidt¹, Axel Fischer¹, Alexander Erban¹, Majken Pagter², Umarah Mubeen¹, Dirk Walther¹, Patrick Giavalisco^{1,3}, Joachim Kopka¹, Heike Sprenger^{1,4}, Dirk K. Hincha¹

¹Max-Planck-Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam, Germany

²Department of Chemistry and Bioscience, Aalborg University, Denmark

³Max Planck Institute for Biology of Aging, Köln, Germany

⁴present address: VIB-UGent Center for Plant Systems Biology, Ghent University, Belgium

Plants from temperate regions increase their freezing tolerance during exposure to low, non-freezing temperatures, a process called cold acclimation or cold priming as plants become better prepared for subsequent stresses. Adaptation to alternating temperatures enables plants to survive winter and to cope with recurring cold periods. In spring loss of freezing tolerance occurs during a tightly regulated deacclimation balancing between transition to reproductive growth and the need for maintained freezing tolerance. Cold priming has been intensively investigated whereas processes underlying cold memory are unknown. We showed that the freezing tolerance of two *Arabidopsis* accessions Col-0 and N14 was higher after priming at 4°C, a lag phase at 20°C and a second cold trigger compared to only primed plants implicating cold stress memory. The triggering response differed from the priming response at the levels of gene expression (RNA-Seq), lipid (UPLC-MS), and metabolite composition (GC-MS). Unique differentially expressed genes after triggering were overrepresented in categories such as lipid and secondary metabolism, stress, redox and cell wall related functions in Col-0 and growth-related functions in N14. Furthermore arabidopsides were accumulated with proposed functions in signalling or as precursors of jasmonic acid. The more freezing tolerant N14 maintained levels of cold-induced metabolites during the lag phase whereas strong accumulation occurred in the less freezing tolerant Col-0 after triggering pointing to genetic differences of transcriptomic and metabolic patterns during cold memory. This is to our knowledge the first report on molecular and metabolic changes accompanying cold stress memory and triggering by a second cold stress.

Session: Abiotic stress and climate change

Subsession: Time, Temperature, and a Transforming World